Grazing muzzles: Do they really work?

×

Sarah Nelson

# Grazing muzzles: Do they really work?

Published August 11, 2016 SHARE

- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Facebook (opens in new window)
- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Twitter (opens in new window)
- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Pinterest (opens in new window)
- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Download (opens in new window)
- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Email (opens in new window)
- <?xml version="1.0" encoding="UTF-8"?> .st0 { fill: #00372b; } Print (opens in same window)

If like many owners you have considered using a grazing muzzle, you may have been left wondering whether or not they really work. Two recent studies have set out to answer this very question!



# **Study One:**

The effect of wearing a grazing muzzle vs. not wearing a grazing muzzle on intakes of spring, summer and autumn pastures by ponies(open in new tab)

### Method

- Grass intake was measured across 3 hours on 4 occasions in 4 ponies with and without a Shires grazing muzzle in spring, summer & autumn
- Grass intake was calculated using changes in body weight (BW), accounting for droppings & urine (which were collected in 'nappies') and natural weight loss due to the evaporation of water in the body

# **Key Results**

- On average grazing muzzles reduced intake by 78% in spring, 77% in summer and 83% in autumn
- Without a muzzle, ponies consumed on average 0.51% BW in spring, 0.57% BW in summer & 0.8% BW in autumn

#### What does this mean?

• Across the seasons, muzzles reduced grass intake by approximately 80% on average and may therefore be an effective method of reducing grass intake throughout the year.



# **Study Two:**

Efficacy of wearing grazing muzzles for 10 hours per day on controlling bodyweight in pastured ponies(open in new tab)

#### Method

- 5 non-obese ponies were turned out for 23 hours per day either without a grazing muzzle (FREEGRAZE treatment) or with a grazing muzzle fitted for 10 out of the 23 hours per day (MUZZLE treatment) during a 6 week crossover study (ponies were divided into 2 groups and were exposed to both treatments)
- Ponies were weighed on alternate days at 8am and 6pm

# **Key Results**

- 4 out 5 ponies gained significantly less weight during the MUZZLE treatment
- 2 ponies exhibited 'pasture damaging' behaviour whilst wearing a muzzle, digging holes of approximately 35cm in diameter with their forefeet
- 1 pony became resentful of wearing the muzzle during week three and lost weight
- 1 pony had to be removed from the study after gaining weight rapidly during the MUZZLE treatment (this pony did not therefore complete the FREEGRAZE treatment)

#### What does this mean?

- Using a grazing muzzle for 10 hours per day may be an effective weight management strategy for some but not all ponies living out at pasture
- Individual horses/ ponies should always be monitored closely for changes in body weight and behaviour

For more advice on how to use grazing muzzles safely call the SPILLERS Care-Line on 01908 226626

@media(min-width:768px){ .freeform-story--left img { margin: 0 auto; width: 35%; } } .article-header-image\_image{ display: none; } @media (min-width: 1024px){ .articleheader-image container{ margin-top: -0.8%; } .freeform-story paragraph{ margin-top: -5%; } } @media (min-width: 767px) and (max-width: 1023px){ .freeformstory\_paragraph{ margin-top: -7%; } } @media (min-width: 275px) and (max-width: 766px){ .freeform-story\_paragraph{ margin-top: -20%; } } .eyebrow.eyebrow--tt-normal{ padding-top: 1%; } .eyebrow{ padding-top:2%; } .card-item{ background: #ffffff; background-image: none !important; } .article-card--with-bg-img .bg-image img { width: 100%; height: auto; max-width: 90%; max-height: 200px; margin-top: 10%; margin-left: 5%; } .article-card-heading{ margin-top: 280px; } .article-card-heading\_link{ font-size: 20px; line-height: 37px; } .article-card-paragraph--xs{ display: none; } .search\_results\_block .search-results-page.card-grid-results .ajax-card-grid\_\_item\_wrapper .article-card-heading link { margin-bottom: 20% !important; font-size: 20px !important; } article-card--with-bg-img .bg-image img{ width: 100%; height: auto; max-width:90%; maxheight:200px; margin-top:10%; } .article-card-heading { margin-top: 212px; } @media (min-width: 1024px) { .article-card .default-link { bottom: 5rem; } .article-card .default-link { margin-left: 21%; } } @media (min-width: 275px) and (max-width: 1023px){ .article-card .default-link { bottom: 6.5rem; } .article-card .default-link { margin-left: 24%; } }

# **Popular Blogs**



# Competing with gastric ulcers: What you need to know

With gastric ulcers thought to affect approximately 60% of competition horses (and up to 50% of leisure horses), it's no surprise that we receive lots of questions about gastric ulcers via our Care-Line.

Read now

### ×

# The importance of lysine for overweight horses

Cutting calories is essential for weight loss but this shouldn't come at the expense of providing a balanced diet. Lysine is an essential amino acid and is one of several nutrients that may be deficient in a calorie restricted diet.

Read now



### Breaking down the diet: What ex-racehorses need to thrive

Whilst the stereotypical ex-racehorse might be prone to excitability, have poor feet and struggle to maintain weight that's certainly not the case for all thoroughbreds. Indeed, once settled into their new workloads and routines, many can be incredibly good-doers!

Read now

×

### **Everything you need to know about laminitis**

With laminitis a worry for many owners we're here to help you brush up on your knowledge of the potential causes, risk factors and management advice!

Read now

# ×

#### What to do when laminitis strikes

Being able to spot the signs of laminitis early maximises your horse's chance of recovery. It pays to be vigilant, especially as subtle signs such as slight reluctance to turn or shortening of stride can be easily missed. While laminitis can strike in any season, a 'flush' of rapid grass growth makes spring a high-risk period, and so can turnout on a sunny frosty morning. Brush up your knowledge of the signs and what to do if you suspect your horse/pony may have laminitis in this blog.

Read now

**Source URL:** https://www.spillers-feeds.com/grazing-muzzles-safety