
Practical ways to reduce forage wastage









Sarah Nelson

Practical ways to reduce forage wastage

Published June 27, 2023

SHARE

-  [Facebook \(opens in new window\)](#)
-  [Twitter \(opens in new window\)](#)
-  [Pinterest \(opens in new window\)](#)
-  [Download \(opens in new window\)](#)
-  [Email \(opens in new window\)](#)
-  [Print \(opens in new window\)](#)

With recent estimations suggesting that almost one third of conserved forage fed to horses ends up being thrown away, we are encouraging horse owners avoid waste by focusing on **how much their horse actually eats** rather than over-supplying forage relative to appetite. However, the premise of feeding as much as a horse will eat (unless they are overweight) is still central to our advice.

```
@media(min-width:768px){ .freeform-story--left img { margin: 0 auto; width: 50%; } }
.article-header-image__image{ display: none; } @media (min-width: 1024px){ .article-
header-image__container{ margin-top: -0.8%; } .freeform-story__paragraph{ margin-top:
-5%; } } @media (min-width: 767px) and (max-width: 1023px){ .freeform-
story__paragraph{ margin-top: -7%; } } @media (min-width: 275px) and (max-width:
766px){ .freeform-story__paragraph{ margin-top: -20%; } } .eyebrow.eyebrow--tt-normal{
padding-top: 1%; } .eyebrow{ padding-top:2%; } .card-item{ background: #ffffff;
```

```
background-image: none !important; } .article-card--with-bg-img .bg-image img { width: 100%; height: auto; max-width: 90%; max-height: 200px; margin-top: 10%; margin-left: 5%; } .article-card-heading { margin-top: 280px; } .article-card-heading__link { font-size: 20px; line-height: 37px; } .article-card-paragraph--xs { display: none; } .search_results_block .search-results-page.card-grid-results .ajax-card-grid__item_wrapper .article-card-heading__link { margin-bottom: 20% !important; font-size: 20px !important; } .article-card--with-bg-img .bg-image img { width: 100%; height: auto; max-width: 90%; max-height: 200px; margin-top: 10%; } .article-card-heading { margin-top: 212px; } @media (min-width: 1024px) { .article-card .default-link { bottom: 5rem; } .article-card .default-link { margin-left: 21%; } } @media (min-width: 275px) and (max-width: 1023px) { .article-card .default-link { bottom: 6.5rem; } .article-card .default-link { margin-left: 24%; } }
```



Wastage

At a recent scientific conference in Belgium (the European Equine Health and Nutrition Congress), there was comprehensive discussion about the amount of forage wasted when feeding horses, with some estimating that in Europe, around 30% of the hay or haylage fed to horses ends up on the muck heap.

“As a company, we are mindful of the environmental pressures we are facing and want to do our part to reduce waste, while still promoting management advice that supports horse welfare,” said Sarah Nelson, Product Manager at Mars Horsecare, home of the SPILLERS™ brand.

Terminology

Traditionally the makers of SPILLERS have used the term ‘ad lib’ when recommending the feeding of forage without restriction. Instead, the team of nutrition advisors will now use language such as ‘ideally horses should be provided with as much forage as they will eat, while being mindful of excess waste.’ An important caveat is that what is actually eaten, not what is given, should always be monitored when feeding conserved forage. It is also important to ensure that an absolute minimum of 1.5% of current bodyweight (on a dry matter basis) of forage is eaten (typically at least 8.5kg of hay for a 500kg horse without access to grazing), unless the horse is on a carefully monitored weight loss diet.

Research

Early research presented at the Equine Science Society Symposium in Texas in June 2023¹ has provided encouraging information on the use of hay nets to prevent waste.

The preliminary study investigated the effect of feeding round bale hay with and without a net on hay usage and dental health. The researchers found that using a net significantly reduced hay wastage without increasing the risk of dental issues and estimated that it could save approximately 1.3 tonnes of wasted hay per year for a 500kg horse.

Haynets/hay boxes

The haybox (which is a large floor-based container with a grill or mesh placed on top of the forage to slow eating time), and other slow feeding devices are also becoming increasingly recognised as an effective way to feed forage safely while minimising waste. These feeding

devices aim to mimic a more natural way of feeding from the ground, while slowing intake.

“While hayboxes and nets may not be suitable for all horses, they are worth considering, both for preventing waste and weight management,” said Sarah. “In fact, as part of our on-going research in the areas of laminitis and obesity, we continue to be involved in several studies investigating the use of hay nets and feeders to help us better understand how we can manage intake without compromising health and welfare.

“We are committed to constantly improving our knowledge and advice in horse nutrition and management. We hope that our change in terminology will help reduce forage waste and have a positive impact on the environment.”

For help estimating forage requirements or for individual advice [contact us](#) on the Care-Line via 01908 226626.

Popular Blogs



[Competing with gastric ulcers: What you need to know](#)

With gastric ulcers thought to affect approximately 60% of competition horses (and up to 50% of leisure horses), it's no surprise that we receive lots of questions about gastric ulcers via our Care-Line.

[Read now](#)



[The importance of lysine for overweight horses](#)

Cutting calories is essential for weight loss but this shouldn't come at the expense of providing a balanced diet. Lysine is an essential amino acid and is one of several nutrients that may be deficient in a calorie restricted diet.

[Read now](#)



[Breaking down the diet: What ex-racehorses need to thrive](#)

Whilst the stereotypical ex-racehorse might be prone to excitability, have poor feet and struggle to maintain weight that's certainly not the case for all thoroughbreds. Indeed, once settled into their new workloads and routines, many can be incredibly good-doers!

[Read now](#)



[Everything you need to know about laminitis](#)

With laminitis season upon us what better time to brush up on your knowledge of the potential causes, risk factors and management advice?

[Read now](#)



What to do when laminitis strikes

Being able to spot the signs of laminitis early maximises your horse's chance of recovery. It pays to be vigilant, especially as subtle signs such as slight reluctance to turn or shortening of stride can be easily missed.

[Read now](#)

Source URL: *<https://www.spillers-feeds.com/moving-away-from-ad-lib-forage>*