






#SPILLERSScience: turnout does not always reduce the risk of gastric ulcers
Danni Twose

#SPILLERSScience: turnout does not always reduce the risk of gastric ulcers

Published June 13, 2023

SHARE

-  [Facebook \(opens in new window\)](#)
-  [Twitter \(opens in new window\)](#)
-  [Pinterest \(opens in new window\)](#)
-  [Download \(opens in new window\)](#)
-  [Email \(opens in new window\)](#)
-  [Print \(opens in new window\)](#)

While the risk of gastric ulcers generally seems lower in horses at pasture, recent research in Iceland, carried out by Nanna Luthersson in collaboration with SPILLERS, the University of Glasgow and the University of Bristol, highlights this may not always be the case. In fact, stabling with suitable forage may even reduce the risk in some situations...

Why Iceland?

Most of the research investigating the prevalence of gastric ulcers has been carried out in 'managed' horses. In Iceland, horses typically live out at pasture, often in large herds and if stabled, are generally fed a high forage, low starch and low sugar diet. While Icelandic horses do get gastric ulcers, it's been suggested that the over-all incidence is low.

✖

```
@media(min-width:768px){ .freeform-story--left img { margin: 0 auto; width: 50%; } }
.article-header-image__image{ display: none; } @media (min-width: 1024px){ .article-
header-image__container{ margin-top: -0.8%; } .freeform-story__paragraph{ margin-top:
-5%; } } @media (min-width: 767px) and (max-width: 1023px){ .freeform-
story__paragraph{ margin-top: -7%; } } @media (min-width: 275px) and (max-width:
766px){ .freeform-story__paragraph{ margin-top: -20%; } } .eyebrow.eyebrow--tt-normal{
padding-top: 1%; } .eyebrow{ padding-top:2%; } .card-item{ background: #ffffff;
background-image: none !important; } .article-card--with-bg-img .bg-image img { width:
100%; height: auto; max-width: 90%; max-height: 200px; margin-top: 10%; margin-left:
5%; } .article-card-heading{ margin-top: 280px; } .article-card-heading__link{ font-size:
20px; line-height: 37px; } .article-card-paragraph--xs{ display: none; }
.search_results_block .search-results-page.card-grid-results .ajax-card-grid__item_wrapper
.article-card-heading__link { margin-bottom: 20% !important; font-size: 20px !important; }
article-card--with-bg-img .bg-image img{ width: 100%; height: auto; max-width:90%; max-
height:200px; margin-top:10%; } .article-card-heading { margin-top: 212px; } @media
(min-width: 1024px){ .article-card .default-link { bottom: 5rem; } .article-card .default-link{
margin-left: 21%; } } @media (min-width: 275px) and (max-width: 1023px){ .article-card
.default-link { bottom: 6.5rem; } .article-card .default-link{ margin-left: 24%; } } table, td {
border: 1px solid #000; text-align: center; border-collapse: collapse; } td { padding: 5px; }
```

What did we do?

The aim of this study was to investigate the incidence of gastric ulcers in Icelandic horses moving from pasture into training.

All horses were scoped within two weeks of being removed from pasture and prior to starting training. They were then scoped again after approximately 8 weeks of being stabled and doing light work.

Prior to the study, all horses were had lived out in large herds for their entire adult lives (age range 3-7 years), had never been in work and were fed supplementary forage in winter months only.

Most horses (57/71) were fed forage only during the training period, but 11 were given very small amounts of soaked sugar beet and 3 were given a small amount of commercially produced feed. Starch and sugar intake from 'bucket feed' in all horses was low - no more than 0.5g per kilogram of bodyweight per day (equivalent to no more than 250g for a 500kg horse).

All the horses in the study were independently owned and although veterinary treatment was offered for all horses if clinically appropriate, it was only accepted for one horse. This horse was later removed from the study and is not included in the results.

What did we find?

- Approximately 72% of horses had non-glandular ulcers (grade 2 or above) at scope 1
- Approximately 25% of horses had non-glandular ulcers (grade 2 or above) at scope 2 – yes that's right, the prevalence and severity of ulcers improved after being stabled for 8 weeks!
- Horses given forage three times per day as opposed to twice per day were more almost 18 times more likely to improve!

-Over-all, the incidence of glandular ulcers decreased from 47% to approximately 41% although it was not possible to identify a reason for this.

	Incidence of non-glandular ulcers at scope 1 (%)	Incidence of non-glandular ulcers at scope 2 (%)
No ulcers present	14	54
Grade 1	15	21
Grade 2	26	14
Grade 3	40	11
Grade 4	6	0

What does this mean?

The high prevalence and severity of non-glandular ulcers at the start of the study, and the subsequent improvement following the training period was unexpected. We are already looking into the reasons for this so watch this space for more information!

If otherwise appropriate, increasing the amount of time your horse is turned out may still help to reduce the risk of gastric ulcers, as well as offering other health and welfare benefits. However, the results of this study highlight the importance of regular forage provision, as well as being a good reminder that 24/7 turnout does not automatically mean reduced risk.



For more advice on feeding horses prone to gastric ulcers contact the [SPILLERS Care-Line](#)

Reference

Luthersson N, Ýr Þorgrímsdóttir Ú, Harris PA, Parkins T, Bennet ED. Effect of moving from being extensively managed out in pasture into training on the incidence of equine gastric ulcer syndrome in Icelandic horses. J Am Vet Med Assoc. 2022 Sep 28;260(S3):S102-S110. doi: 10.2460/javma.22.06.0263. PMID: 36149938.

Popular Blogs



[Competing with gastric ulcers: What you need to know](#)

With gastric ulcers thought to affect approximately 60% of competition horses (and up to 50% of leisure horses), it's no surprise that we receive lots of questions about gastric ulcers via our Care-Line.

[Read now](#)



The importance of lysine for overweight horses

Cutting calories is essential for weight loss but this shouldn't come at the expense of providing a balanced diet. Lysine is an essential amino acid and is one of several nutrients that may be deficient in a calorie restricted diet.

[Read now](#)



Breaking down the diet: What ex-racehorses need to thrive

Whilst the stereotypical ex-racehorse might be prone to excitability, have poor feet and struggle to maintain weight that's certainly not the case for all thoroughbreds. Indeed, once settled into their new workloads and routines, many can be incredibly good-doers!

[Read now](#)



Everything you need to know about laminitis

With laminitis season upon us what better time to brush up on your knowledge of the potential causes, risk factors and management advice?

[Read now](#)



What to do when laminitis strikes

Being able to spot the signs of laminitis early maximises your horse's chance of recovery. It pays to be vigilant, especially as subtle signs such as slight reluctance to turn or shortening of stride can be easily missed.

[Read now](#)

Source URL:

<https://www.spillers-feeds.com/spillersscience-turnout-does-not-always-reduce-the-risk-of-gastric-ulcers>