
#SPILLERSScience: Reasons for retirement

#SPILLERSScience: Reasons for retirement

SHARE

- [Facebook \(opens in new window\)](#)
- [Twitter \(opens in new window\)](#)
- [Pinterest \(opens in new window\)](#)
- [Download \(opens in new window\)](#)
- [<?xml version="1.0" encoding="UTF-8"?> Email \(opens in new window\)](mailto:SPILLERS@uky.edu)
- [Print \(opens in new window\)](#)



Survey results recently published by the Gluck Equine Research Center at the University of Kentucky in partnership with SPILLERS have shed light on the reasons why horses may be retired.

By developing a better understanding of when and why senior horses are being retired we hope, eventually, to be better placed to monitor for, prevent (where possible), and treat the underlying conditions early on. The overall aim being to try to help horses stay active and healthy for longer.

Survey basics

A total of 2717 surveys completed by horse owners in the US were analysed. Approximately 65% of respondents were leisure owners and approximately 35% were equine professionals. Most of the horses were being used for pleasure riding or driving (~39%) or were in full retirement (~40%).

Age & health

Most horses (62%) were retired between 15-24 years of age, with health problems given as the main reasons. Horses with owner reported veterinary-diagnosed laminitis, lameness and Degenerative suspensory ligament desmitis had higher odds of retirement than those that did not have these conditions.

Mares vs geldings

Mares had a higher risk of retirement than geldings. This may be partially due to broodmares experiencing a reduction or absence of regular exercise.

Muscle matters

Around 1 in 6 horses had low muscle mass (muscle wastage) according to their owners and in most cases, owners felt this affected their horse's welfare and ability to work.

- Age, sex, osteoarthritis, laminitis, pituitary pars intermedia dysfunction (PPID), and primary use were identified as risk factors for low muscle mass.
- Geldings had a higher risk of low muscle mass than mares, which may be a result of low testosterone.
- Horses with owner-reported veterinary-diagnosed PPID (also known as 'Cushing's syndrome'), osteoarthritis and laminitis had a higher risk of low muscle mass.
- Retired horses were at higher risk of low muscle mass than horses used for competition or pleasure riding.

What's next?

In follow-up to this research, we have been looking at how senior horses in the US are managed and their health status. Watch this space!

Further information

This research has been accepted for publication in the Equine Veterinary Journal. You can read the abstract (research summary) [here](#)

We are proud to have been working with the University of Kentucky on a number of senior research projects since 2013. You can learn more about the University of Kentucky's Aged Horse Research Center [here](#)

Reference

Herbst A, Coleman M, Macon E, Brokman A, Stromberg A, Harris P, Adams A (2023) Retirement risk factors, exercise management, and muscle mass in U.S. senior horses Equine Veterinary Journal in press.

```
@media(min-width:768px){ .freeform-story--left img { margin: 0 auto; width: 50%; } }
.article-header-image__image{ display: none; } @media (min-width: 1024px){ .article-
header-image__container{ margin-top: -0.8%; } .freeform-story__paragraph{ margin-top:
-5%; } } @media (min-width: 767px) and (max-width: 1023px){ .freeform-
story__paragraph{ margin-top: -7%; } } @media (min-width: 275px) and (max-width:
766px){ .freeform-story__paragraph{ margin-top: -20%; } } .eyebrow.eyebrow--tt-normal{
padding-top: 1%; } .eyebrow{ padding-top:2%; } .card-item{ background: #ffffff;
background-image: none !important; } .article-card--with-bg-img .bg-image img { width:
100%; height: auto; max-width: 90%; max-height: 200px; margin-top: 10%; margin-left:
5%; } .article-card-heading{ margin-top: 280px; } .article-card-heading__link{ font-size:
20px; line-height: 37px; } .article-card-paragraph--xs{ display: none; }
.search_results_block .search-results-page.card-grid-results .ajax-card-grid__item_wrapper
.article-card-heading__link { margin-bottom: 20% !important; font-size: 20px !important; }
article-card--with-bg-img .bg-image img{ width: 100%; height: auto; max-width:90%; max-
height:200px; margin-top:10%; } .article-card-heading { margin-top: 212px; } @media
(min-width: 1024px){ .article-card .default-link { bottom: 5rem; } .article-card .default-link{
margin-left: 21%; } } @media (min-width: 275px) and (max-width: 1023px){ .article-card
.default-link { bottom: 6.5rem; } .article-card .default-link{ margin-left: 24%; } }
```

Popular Blogs



[Competing with gastric ulcers: What you need to know](#)

With gastric ulcers thought to affect approximately 60% of competition horses (and up to 50% of leisure horses), it's no surprise that we receive lots of questions about gastric ulcers via our Care-Line.

[Read now](#)



The importance of lysine for overweight horses

Cutting calories is essential for weight loss but this shouldn't come at the expense of providing a balanced diet. Lysine is an essential amino acid and is one of several nutrients that may be deficient in a calorie restricted diet.

[Read now](#)



Breaking down the diet: What ex-racehorses need to thrive

Whilst the stereotypical ex-racehorse might be prone to excitability, have poor feet and struggle to maintain weight that's certainly not the case for all thoroughbreds. Indeed, once settled into their new workloads and routines, many can be incredibly good-doers!

[Read now](#)



Everything you need to know about laminitis

With laminitis season upon us what better time to brush up on your knowledge of the potential causes, risk factors and management advice?

[Read now](#)



What to do when laminitis strikes

Being able to spot the signs of laminitis early maximises your horse's chance of recovery. It pays to be vigilant, especially as subtle signs such as slight reluctance to turn or shortening of stride can be easily missed.

[Read now](#)

Source URL: <https://www.spillers-feeds.com/articles/spillersscience-reasons-retirement>